MEDICATION DEFERRAL LIST

Please tell us if you are now taking or if you have EVER taken any of these medications:

Proscar© (finasteride) – usually given for prostate gland enlargement
Avodart©, Jalyn© (dutasteride) – usually given for prostate enlargement
Propecia© (finasteride) – usually given for baldness
Accutane© (Amnesteem, Claravis, Sotret, isotretinoin) – usually given for severe acne
Soriatane© (acitretin) - usually given for severe psoriasis
Tegison© (etretinate) – usually given for severe psoriasis
Growth Hormone from Human Pituitary Glands – used usually for children with delayed or impaired growth
Insulin from Cows (Bovine, or Beef, Insulin) – used to treat diabetes
Hepatitis B Immune Globulin – given following an exposure to hepatitis B. NOTE: This is different from the hepatitis B vaccine which is a series of 3 injections given over a 6 month period to prevent future infection from exposures to hepatitis B.
Unlicensed Vaccine – usually associated with a research protocol

IF YOU WOULD LIKE TO KNOW WHY THESE MEDICINES AFFECT YOU AS A BLOOD DONOR, PLEASE KEEP READING:

- If you have taken or are taking <u>Proscar, Avodart, Propecia, Accutane, Soriatane, or Tegison</u>, these medications can cause birth defects. Your donated blood could contain high enough levels to damage the unborn baby if transfused to a pregnant woman. Once the medication has been cleared from your blood, you may donate again. Following the last dose, the deferral period is one month Proscar, Propecia and Accutane, six months for Avodart and three years for Soriatane. Tegison is a permanent deferral.
- <u>Growth hormone from human pituitary glands</u> was prescribed for children with delayed or impaired growth. The hormone was obtained from human pituitary glands, which are found in the brain. Some people who took this hormone developed a rare nervous system condition called Creutzfeldt-Jakob Disease (CJD, for short). The deferral is permanent.
- Insulin from cows (bovine, or beef, insulin) is an injected material used to treat diabetes. If this insulin was imported into the US from countries in which "Mad Cow Disease" has been found, it could contain material from infected cattle. There is concern that "Mad Cow Disease" is transmitted by transfusion. The deferral is indefinite.
- Hepatitis B Immune Globulin (HBIG) is an injected material used to prevent infection following an exposure to
 hepatitis B. HBIG does not prevent hepatitis B infection in every case, therefore persons who have received HBIG
 must wait 12 months to donate blood to be sure they were not infected since hepatitis B can be transmitted through
 transfusion to a patient.
- <u>Unlicensed Vaccine</u> is usually associated with a research protocol and the effect on blood transmission is unknown. Deferral is one year unless otherwise indicated by Medical Director..